

Discussion Guide

You may have questions about acute myeloid leukemia (AML) and your treatment with VYXEOS (vix-e-ose) that you want to discuss with your doctor and healthcare team.

This discussion guide is designed to help you get the conversation started and ensure that you are comfortable discussing all aspects of your treatment.

Consider the following questions before your next check-in with your healthcare team:



How are you feeling since your last check-in?

How have you been dealing with your diagnosis?

What concerns you most about your diagnosis and treatment?

Do you have any specific questions for your doctor?

WHAT IS VYXEOS?

VYXEOS is an intravenous (IV) chemotherapy used for the treatment of certain types of newly-diagnosed acute myeloid leukemia (AML) in adults and pediatric patients 1 year and older, including patients whose AML is related to previously received chemotherapy or radiation therapy (also called therapy-related AML, t-AML) and AML in patients who previously had certain types of blood disorders (also called AML with myelodysplasia-related changes, AML-MRC).

IMPORTANT SAFETY INFORMATION

WARNING: VYXEOS has different dosage recommendations from other medications that contain daunorubicin and/or cytarabine. Do not substitute VYXEOS for other daunorubicin and/or cytarabine-containing products.

VYXEOS should not be given to patients who have a history of serious allergic reaction to daunorubicin, cytarabine, or any of its ingredients.

Please see additional Important Safety Information throughout and full Prescribing Information, including BOXED Warning, and discuss with your doctor.

Vyxeos[®]
 (daunorubicin and cytarabine) 44 mg/100 mg
 per vial
 liposome for injection

Consider reviewing the following questions with your healthcare team:



How does my medical history affect my diagnosis and my treatment with VYXEOS?

Why is VYXEOS an appropriate treatment for me?

What does the timeline of my treatment look like as of right now?

Where will I receive my treatment?

IMPORTANT SAFETY INFORMATION, continued

VYXEOS can cause a severe decrease in blood cells (red and white blood cells and cells that prevent bleeding, called platelets) which can result in serious infection or bleeding and possibly lead to death. Your doctor will monitor your blood counts during treatment with VYXEOS. Patients should tell the doctor about new onset fever or symptoms of infection or if they notice signs of bruising or bleeding.

VYXEOS can cause heart-related side effects. Tell your doctor about any history of heart disease, radiation to the chest, or previous chemotherapy. Inform your doctor if you develop symptoms of heart failure such as:

- shortness of breath or trouble breathing
- swelling or fluid retention, especially in the feet, ankles, or legs
- unusual tiredness

Please see additional Important Safety Information throughout and full Prescribing Information, including BOXED Warning, and discuss with your doctor.



Consider reviewing the following questions with your healthcare team:



What are the possible side effects of treatment with VYXEOS?

How will we be able to tell if my treatment is working?

How can I prepare for my treatment with VYXEOS?

Where can I find a support community of others who have AML or related conditions?

IMPORTANT SAFETY INFORMATION, continued

VYXEOS may cause allergic reactions including anaphylaxis. Seek immediate medical attention if you develop signs and symptoms of anaphylaxis such as:

- trouble breathing
- severe itching
- skin rash or hives
- swelling of the face, lips, mouth, or tongue

Please see additional Important Safety Information throughout and full Prescribing Information, including BOXED Warning, and discuss with your doctor.



Consider reviewing the following questions with your healthcare team:



What kind of diet and exercise regimen do you recommend for me during treatment?

Do you have any guidance for my caregivers, family, and friends throughout this process?

IMPORTANT SAFETY INFORMATION, continued

VYXEOS contains copper and may cause copper overload in patients with Wilson's disease or other copper-processing disorders.

VYXEOS can damage the skin if it leaks out of the vein. Tell your doctor right away if you experience symptoms of burning, stinging, or blisters and skin sores at the injection site.

VYXEOS can harm your unborn baby. Inform your doctor if you are pregnant, planning to become pregnant, or nursing. Do not breastfeed while receiving VYXEOS. Females and males of reproductive potential should use effective contraception during treatment and for 6 months following the last dose of VYXEOS.

The most common side effects were bleeding events, fever, rash, swelling, nausea, sores in the mouth or throat, diarrhea, constipation, muscle pain, tiredness, stomach pain, difficulty breathing, headache, cough, decreased appetite, irregular heartbeat, pneumonia, blood infection, chills, sleep disorders, and vomiting.

Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report side effects to Jazz Pharmaceuticals at 1-800-520-5568.

Please see additional Important Safety Information throughout and full Prescribing Information, including BOXED Warning, and discuss with your doctor.